

# PARENT STOCK

ROSS 308

Nutrition  
Specifications

2016



## Introduction

This booklet contains the nutritional recommendations for Ross® 308 (slow feathering) parent stock and is to be used with the **Ross Parent Stock Management Handbook** and the Ross 308 Parent Stock Performance Objectives.

## Performance

To achieve optimal reproductive performance, it is important that the body-weight profiles recommended in the Ross 308 Parent Stock Performance Objectives are followed. For the nutritional recommendations that follow, nutrient specifications presented have been based upon daily energy allocations that enable body-weight profiles to be achieved.

Included in this booklet are 3 different rearing programs applicable for the following situations:

- **2-Stage Rearing Program** - where the same energy density is used during rear.
- **3-Stage Rearing Program** - where a Pre-Breeder diet is introduced to prepare the birds for the laying period.
- **4-Stage Rearing Program** - where a lower nutrient density and a higher feed volume feeding strategy is required.

**Please note, these nutrient recommendations are based on the specified energy levels. Adjustment of nutrient levels must be made to reflect the feeding of different energy levels. Feed allocation should be determined by body weight and egg production levels, and therefore altered to maintain the recommended weight and egg production profiles.**

It may be beneficial to use a separate male diet during the production period. A specification for a male diet is provided in this booklet.

For further information regarding these recommendations or for more specialized situations and advice on local markets, please contact your Aviagen® Nutritionist or any Aviagen representative.

## Contents

03	2-Stage Rearing Program
04	3-Stage Rearing Program
05	4-Stage Rearing Program
06	Male Program

## Female Parent Stock Nutrient Specifications

### 2-Stage Rearing Program

		Starter		Grower		Breeder 1		Breeder 2 <sup>a</sup>		Breeder 3 <sup>a</sup>	
Age Fed	days	0-28 days		29 days to 5% production		5% production to 245 days		246-350 days		After 351 days	
Energy per kg	kcal	2800		2800		2800		2800		2800	
	MJ	11.70		11.70		11.70		11.70		11.70	
Energy per lb	kcal	1270		1270		1270		1270		1270	
<b>AMINO ACIDS*</b>		<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>
Lysine	%	1.06	0.95	0.68	0.61	0.67	0.60	0.62	0.56	0.58	0.52
Methionine + Cystine	%	0.84	0.74	0.63	0.55	0.67	0.59	0.65	0.57	0.59	0.54
Methionine	%	0.51	0.46	0.38	0.35	0.41	0.37	0.40	0.36	0.36	0.35
Threonine	%	0.75	0.66	0.54	0.48	0.55	0.49	0.53	0.47	0.51	0.47
Valine	%	0.80	0.71	0.64	0.57	0.63	0.56	0.60	0.53	0.57	0.51
IsoLeucine	%	0.70	0.62	0.56	0.50	0.56	0.50	0.54	0.48	0.51	0.45
Arginine	%	1.17	1.05	0.84	0.76	0.88	0.79	0.86	0.77	0.80	0.72
Tryptophan	%	0.19	0.16	0.16	0.14	0.16	0.14	0.15	0.13	0.14	0.12
Leucine	%	1.23	1.11	0.84	0.76	1.04	0.94	1.00	0.90	0.96	0.86
Crude Protein	%	19.00		14.00-15.00		15.00		14.00		13.00	
<b>MINERALS*</b>											
Calcium	%	1.00		0.90		3.00		3.20		3.40	
Available Phosphorus	%	0.45		0.42		0.35		0.33		0.32	
Sodium	%	0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23	
Chloride	%	0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23	
Potassium	%	0.40-0.90		0.40-0.90		0.60-0.90		0.60-0.90		0.60-0.90	
<b>ADDED TRACE MINERALS PER KG</b>											
Copper	mg	16						10			
Iodine	mg	1.25						2.00			
Iron	mg	40						50			
Manganese	mg	120						120			
Selenium	mg	0.30						0.30			
Zinc	mg	110						110			
<b>ADDED VITAMINS PER KG</b>											
		<b>Wheat based feed</b>		<b>Maize based feed</b>				<b>Wheat based feed</b>		<b>Maize based feed</b>	
Vitamin A	IU	11000		10000				12000		11000	
Vitamin D3	IU	3500		3500				3500		3500	
Vitamin E	IU	100		100				100		100	
Vitamin K (Menadione)	mg	3		3				5		5	
Thiamin (B1)	mg	3		3				3		3	
Riboflavin (B2)	mg	6		6				12		12	
Nicotinic Acid	mg	30		35				50		55	
Pantothenic Acid	mg	13		15				13		15	
Pyridoxine (B6)	mg	4		3				5		4	
Biotin	mg	0.20		0.15				0.30		0.25	
Folic Acid	mg	1.50		1.50				2.00		2.00	
Vitamin B12	mg	0.02		0.02				0.03		0.03	
<b>MINIMUM SPECIFICATION</b>											
Choline per kg	mg	1400		1300		1200		1050		1050	
Linoleic Acid	%	1.00		1.00		1.25		1.25		1.25	

Digest = Digestible

\* Energy base value. Nutrients should be factored accordingly when feeding differing energy values.

<sup>a</sup> Breeder 2 and 3 can be useful to help control egg size and improve shell quality.

**NOTES** - These feed specifications should be used as a guide. They may require adjustment for local conditions, legislation and markets.

## Female Parent Stock Nutrient Specifications

### 3-Stage Rearing Program

		Starter		Grower		Pre-Breeder		Breeder 1		Breeder 2 <sup>a</sup>		Breeder 3 <sup>a</sup>	
Age Fed	days	0-28 days		29-133 days		134 days to 5% production		5% production to 245 days		246-350 days		After 351 days	
Energy per kg	kcal	2800		2800		2800		2800		2800		2800	
	MJ	11.70		11.70		11.70		11.70		11.70		11.70	
Energy per lb	kcal	1270		1270		1270		1270		1270		1270	
<b>AMINO ACIDS*</b>		<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>
Lysine	%	1.06	0.95	0.68	0.61	0.60	0.54	0.67	0.60	0.62	0.56	0.58	0.52
Methionine + Cystine	%	0.84	0.74	0.63	0.55	0.59	0.52	0.67	0.59	0.65	0.57	0.59	0.54
Methionine	%	0.51	0.46	0.38	0.35	0.36	0.33	0.41	0.37	0.40	0.36	0.36	0.35
Threonine	%	0.75	0.66	0.54	0.48	0.49	0.43	0.55	0.49	0.53	0.47	0.51	0.47
Valine	%	0.80	0.71	0.64	0.57	0.53	0.47	0.63	0.56	0.60	0.53	0.57	0.51
IsoLeucine	%	0.70	0.62	0.56	0.50	0.48	0.43	0.56	0.50	0.54	0.48	0.51	0.45
Arginine	%	1.17	1.05	0.84	0.76	0.77	0.69	0.88	0.79	0.86	0.77	0.80	0.72
Tryptophan	%	0.19	0.16	0.16	0.14	0.15	0.13	0.16	0.14	0.15	0.13	0.14	0.12
Leucine	%	1.23	1.11	0.84	0.76	0.83	0.75	1.04	0.94	1.00	0.90	0.96	0.86
Crude Protein	%	19.00		14.00-15.00		14.50		15.00		14.00		13.00	
<b>MINERALS*</b>													
Calcium	%	1.00		0.90		1.20		3.00		3.20		3.40	
Available Phosphorus	%	0.45		0.42		0.35		0.35		0.33		0.32	
Sodium	%	0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23	
Chloride	%	0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23	
Potassium	%	0.40-0.90		0.40-0.90		0.60-0.90		0.60-0.90		0.60-0.90		0.60-0.90	
<b>ADDED TRACE MINERALS PER KG</b>													
Copper	mg			16						10			
Iodine	mg			1.25						2.00			
Iron	mg			40						50			
Manganese	mg			120						120			
Selenium	mg			0.30						0.30			
Zinc	mg			110						110			
<b>ADDED VITAMINS PER KG</b>				<b>Wheat based feed</b>	<b>Maize based feed</b>			<b>Wheat based feed</b>	<b>Maize based feed</b>				
Vitamin A	IU			11000	10000			12000	11000				
Vitamin D3	IU			3500	3500			3500	3500				
Vitamin E	IU			100	100			100	100				
Vitamin K (Menadione)	mg			3	3			5	5				
Thiamin (B1)	mg			3	3			3	3				
Riboflavin (B2)	mg			6	6			12	12				
Nicotinic Acid	mg			30	35			50	55				
Pantothenic Acid	mg			13	15			13	15				
Pyridoxine (B6)	mg			4	3			5	4				
Biotin	mg			0.20	0.15			0.30	0.25				
Folic Acid	mg			1.50	1.50			2.00	2.00				
Vitamin B12	mg			0.02	0.02			0.03	0.03				
<b>MINIMUM SPECIFICATION</b>													
Choline per kg	mg	1400		1300		1200		1200		1050		1050	
Linoleic Acid	%	1.00		1.00		1.00		1.25		1.25		1.25	

Digest = Digestible

\* Energy base value. Nutrients should be factored accordingly when feeding differing energy values.

<sup>a</sup> Breeder 2 and 3 can be useful to help control egg size and improve shell quality.

**NOTES** - These feed specifications should be used as a guide. They may require adjustment for local conditions, legislation and markets.

## Female Parent Stock Nutrient Specifications

### 4-Stage Rearing Program

		Starter 1		Starter 2		Grower		Pre-Breeder		Breeder 1		Breeder 2 <sup>a</sup>		Breeder 3 <sup>a</sup>	
Age Fed	days	0-21 days		22-35 days		36-105 days		106 days to 5% production		5% production to 245 days		246-350 days		After 351 days	
Energy per kg	kcal	2800		2800		2600		2700		2800		2800		2800	
	MJ	11.70		11.70		10.90		11.30		11.70		11.70		11.70	
Energy per lb	kcal	1270		1270		1179		1225		1270		1270		1270	
<b>AMINO ACIDS*</b>		<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>	<b>Total</b>	<b>Digest</b>
Lysine	%	1.06	0.95	0.74	0.67	0.58	0.52	0.58	0.52	0.67	0.60	0.62	0.56	0.58	0.52
Meth + Cyst	%	0.84	0.74	0.67	0.59	0.59	0.52	0.58	0.51	0.67	0.59	0.65	0.57	0.59	0.54
Methionine	%	0.51	0.46	0.41	0.37	0.36	0.33	0.35	0.32	0.41	0.37	0.40	0.36	0.36	0.35
Threonine	%	0.75	0.66	0.60	0.53	0.50	0.44	0.47	0.41	0.55	0.49	0.53	0.47	0.51	0.47
Valine	%	0.80	0.71	0.70	0.63	0.49	0.44	0.51	0.45	0.63	0.56	0.60	0.53	0.57	0.51
IsoLeucine	%	0.70	0.62	0.62	0.55	0.45	0.40	0.47	0.41	0.56	0.50	0.54	0.48	0.51	0.45
Arginine	%	1.17	1.05	0.93	0.83	0.71	0.64	0.74	0.67	0.88	0.79	0.86	0.77	0.80	0.72
Tryptophan	%	0.19	0.16	0.18	0.15	0.14	0.12	0.15	0.13	0.16	0.14	0.15	0.13	0.14	0.12
Leucine	%	1.23	1.11	0.93	0.83	0.77	0.69	0.80	0.72	1.04	0.94	1.00	0.90	0.96	0.86
Crude Protein	%	19.00		17.00		13.00-14.00		14.00		15.00		14.00		13.00	
<b>MINERALS*</b>															
Calcium	%	1.00		1.00		0.90		1.20		3.00		3.20		3.40	
Av. Phosphorus	%	0.45		0.45		0.42		0.35		0.35		0.33		0.32	
Sodium	%	0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23	
Chloride	%	0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23		0.18-0.23	
Potassium	%	0.40-0.90		0.40-0.90		0.40-0.90		0.60-0.90		0.60-0.90		0.60-0.90		0.60-0.90	
<b>ADDED TRACE MINERALS PER KG</b>															
Copper	mg			16								10			
Iodine	mg			1.25								2.00			
Iron	mg			40								50			
Manganese	mg			120								120			
Selenium	mg			0.30								0.30			
Zinc	mg			110								110			
<b>ADDED VITAMINS PER KG</b>															
Vitamin A	IU			11000		10000						12000		11000	
Vitamin D3	IU			3500		3500						3500		3500	
Vitamin E	IU			100		100						100		100	
Vitamin K	mg			3		3						5		5	
Thiamin (B1)	mg			3		3						3		3	
Riboflavin (B2)	mg			6		6						12		12	
Nicotinic Acid	mg			30		35						50		55	
Pantothenic Acid	mg			13		15						13		15	
Pyridoxine (B6)	mg			4		3						5		4	
Biotin	mg			0.20		0.15						0.30		0.25	
Folic Acid	mg			1.50		1.50						2.00		2.00	
Vitamin B12	mg			0.02		0.02						0.03		0.03	
<b>MINIMUM SPECIFICATION</b>															
Choline per kg	mg	1400		1400		1300		1200		1200		1050		1050	
Linoleic Acid	%	1.00		1.00		1.00		1.00		1.25		1.25		1.25	

Digest = Digestible

\* Energy base value. Nutrients should be factored accordingly when feeding differing energy values.

<sup>a</sup> Breeder 2 and 3 can be useful to help control egg size and improve shell quality.

**NOTES** - These feed specifications should be used as a guide. They may require adjustment for local conditions, legislation and markets.

## Male Parent Stock Nutrient Specifications

Feed allocation will be determined by male body weight and condition.  
The male diet should be introduced when birds are moved to the laying house or at light stimulation.

		Male Feed	
Energy per kg	kcal	2700	
	MJ	11.30	
Energy per lb	kcal	1225	
AMINO ACIDS*		Total	Digest
Lysine	%	0.49	0.44
Methionine + Cystine	%	0.48	0.42
Methionine	%	0.31	0.28
Threonine	%	0.38	0.33
Valine	%	0.42	0.37
IsoLeucine	%	0.39	0.34
Arginine	%	0.58	0.52
Tryptophan	%	0.09	0.08
Leucine	%	0.58	0.52
Crude Protein	%	11.50	
MINERALS*			
Calcium	%	0.70	
Available Phosphorus	%	0.35	
Sodium	%	0.18-0.23	
Chloride	%	0.18-0.23	
Potassium	%	0.60-0.90	
ADDED TRACE MINERALS PER KG			
Copper	mg	10	
Iodine	mg	2.00	
Iron	mg	50	
Manganese	mg	120	
Selenium	mg	0.30	
Zinc	mg	110	
ADDED VITAMINS PER KG		Wheat based feed	Maize based feed
Vitamin A	IU	12000	11000
Vitamin D3	IU	3500	3500
Vitamin E	IU	100	100
Vitamin K (Menadione)	mg	5	5
Thiamin (B1)	mg	3	3
Riboflavin (B2)	mg	12	12
Nicotinic Acid	mg	50	55
Pantothenic Acid	mg	13	15
Pyridoxine (B6)	mg	5	4
Biotin	mg	0.30	0.25
Folic Acid	mg	2.00	2.00
Vitamin B12	mg	0.03	0.03
MINIMUM SPECIFICATION			
Choline per kg	mg	1000	
Linoleic Acid	%	1.00	

Digest = Digestible

\* Energy base value. Nutrients should be factored accordingly when feeding differing energy values.

**NOTES** - These feed specifications should be used as a guide. They may require adjustment for local conditions, legislation and markets.





[www.aviagen.com](http://www.aviagen.com)

Every attempt has been made to ensure the accuracy and relevance of the information presented. However, Aviagen accepts no liability for the consequences of using the information for the management of chickens.

For further information, please contact your local Aviagen representative.

Aviagen and the Aviagen logo, and Ross and the Ross logo are registered trademarks of Aviagen in the US and other countries. All other trademarks or brands are registered by their respective owners.